arXiv:1612.01348v1 [math.DG] 5 Dec 2016

THE CONTINUITY METHOD ON FANO BUNDLES

YASHAN ZHANG AND ZHENLEI ZHANG

Abstract. We prove that the continuity method on a Fano bundle starting from a suitable Kähler metric converges to a Kähler metric on the base in Gromov-Hausdorff topology.

1. Motivation and main result

In [16], La Nave and Tian introduced a new approach to the Analytic Minimal Model Program. It is a continuity method of complex Monge-Ampère equations.

As we know from birational geometry, Fano fibration is considered to be a standard form for projective variety which is not birational to a minimal model. In fact, Fano fibration arises naturally as one type of the extremal contractions (see [7, 17] for details and examples). Therefore, to carry out Analytic Minimal Model Program, it should be natural to study the behaviours of the continuity method on Fano fibrations.

In this paper, motivated by above, we will study the geometric convergence of the continuity method on Fano bundles. To state our main result, we now fix some notations.

Let X be an n-dimensional compact connected Kähler manifold admitting a Fano bundle over a k-dimensional compact connected Kähler manifold B with $k < n$. Namely, there exists a surjective holomorphic submersion $f : X \to B$ with connected fibers such that $K_X^{-1}|_{f^{-1}(b)}$ is ample for all $b \in B$. Set $X_b := f^{-1}(b)$, which are $(n-k)$-dimensional Fano manifolds since $K_X^{-1}|_{X_b} \cong K_{X_b}^{-1}$ by adjunction formula. According to Lemma 2.2 in [31], we fix a Kähler class α on X and a Kähler class β on B such that $\alpha = 2\pi c_1(X) + f^*\beta$.

For any Kähler metric $\omega_0 \in \alpha$, consider the following continuity method introduced by La Nave and Tian in [16] and Rubinstein in [18]:

$$\begin{cases} \omega(t) = \omega_0 - tRic(\omega(t)) \\ \omega(0) = \omega_0. \end{cases} \tag{1.1}$$

It is proved in [16] that the maximal existence time of (1.1) is

$$T = \sup\{t > 0 | [\omega_0] - 2\pi t c_1(X) > 0\}.$$

It is not hard to see that $T = 1$ in our case and the limiting class satisfies

$$[\omega_0] - 2\pi c_1(X) = f^*\beta, \tag{1.2}$$

which in particular implies

$$\omega_0|_{X_b} \in 2\pi c_1(X_b). \tag{1.3}$$

We should remark that, in general, if we choose $\tilde{\alpha}$ and $\tilde{\beta}$ with $\tilde{\alpha} = 2\pi t_0 c_1(X) + f^*\tilde{\beta}$ for some positive constant t_0, then the maximal existence time $T = t_0$ and all our discussions apply with mild modifications.

Y. Zhang is partially supported by the Science and Technology Development Fund (Macao S.A.R.) Grant FDCT/ 016/2013/A1 and the Project MYRG2015-00235-FST of the University of Macau.

Z. Zhang is partially supported by NSFC 11431009.

According to Conjecture 4.1 of [16], $(X, \omega(t))$ should converge to a metric d_B on B in Gromov-Hausdorff topology as $t \to 1$. In this paper we shall confirm this conjecture in above situation. More precisely, we will prove the following

Theorem 1.1. *For any Kähler metric $\omega_0 \in \alpha$, there exists a Kähler metric $\bar{\omega}_B \in \beta$ such that the unique solution $\omega(t)$ of (1.1) starting from ω_0 converges to $(B, \bar{\omega}_B)$ in Gromov-Hausdorff topology as $t \to 1$.*

Let's discuss some special cases.

Example 1.2. *If $k = 0$ (i.e., B is a point) and hence X is a Fano manifold, then by Theorem 1.1 the continuity method (1.1) starting from any Kähler metric $\omega_0 \in 2\pi c_1(X)$ collapses to a point in Gromov-Hausdorff topology. In fact, in this case it is natural to consider a normalized version of (1.1) as follows*

$$(1-t)\omega(t) = \omega_0 - tRic(\omega(t)). \tag{1.4}$$

By Theorem 1.1 of [16] (1.4) has a unique smooth solution $\omega(t)$ for $t \in [0, 1)$. Moreover, by using some well-known results in complex Monge-Ampère equations (see Section 3.1 of [6] for detailed arguments) we have that, as $t \to 1^-$, $\omega(t)$ converges in $C^\infty(X, \omega_0)$-topology to a unique Kähler metric $\omega(1) \in 2\pi c_1(X)$ satisfying $Ric(\omega(1)) = \omega_0$ (this observation motivates some discussions in Section 4, see Remark 4.3 and Proposition 4.4). Then the time t will go through 1 and (1.4) can be written as

$$Ric(\omega(t)) = \frac{1}{t}\omega_0 + (1 - \frac{1}{t})\omega(t), \tag{1.5}$$

which is a simple reparametrization of the classical continuity method proposed by Aubin [1, 2]. Therefore, all the results concerning Aubin's continuity method on Fano manifolds (see, e.g., [8] for recent progress) apply in our case. For example, by the results in [3], if X admits a unique Kähler-Einstein metric $\omega_{KE} \in 2\pi c_1(X)$, then (1.4) (or (1.5)) is solvable for $t \in [1, \infty)$ and $\omega(t)$ converges in $C^\infty(X, \omega_0)$-topology to ω_{KE} as $t \to \infty$.

The next example will fit the motivation and Theorem 1.1 well.

Example 1.3. *Let Σ_a, $a \in \mathbb{Z}_{\geq 1}$, be a complete smooth fan in $\mathbb{R}^2$ with the minimal generators $u_0 = (-1, a)$, $u_1 = (0, 1)$, $u_2 = (1, 0)$ and $u_3 = (0, -1)$. Let $\mathcal{H}_a$ be the smooth toric variety corresponding to Σ_a, which is called the a-th Hirzebruch surface. We shall explain how to equip a $\mathbb{CP}^1$-bundle structure on $\mathcal{H}_a$ by extremal contractions. To this end, firstly note that the wall $\tau = \mathbb{R}_{\geq 0} \cdot u_2$ is an extremal wall and will give an extremal ray $\mathcal{R}$ in the Mori cone of $\mathcal{H}_a$ (see Example 6.3.23 in [7]). Moreover, since the wall relation of τ is*

$$u_1 + 0 \cdot u_2 + u_3 = 0,$$

by Proposition 15.4.5 of [7] the extremal contraction given by $\mathcal{R}$ is a fibration (i.e., Mori fiber space) $f : \mathcal{H}_a \to \mathbb{CP}^1$ and the fibers are isomorphic to the toric variety of the complete fan in $\mathbb{R}$, that is $\mathbb{CP}^1$. Hence all fibers are smooth and f is in fact a locally trivial fibration (see p.190 of [4]). Consequently, f is a $\mathbb{CP}^1$-bundle. Now the discussions in Section V.4 (in particular, Proposition V.4.2) of [4] imply that f is exactly the projective bundle $\mathbb{P}(\mathcal{O}_{\mathbb{CP}^1} \otimes \mathcal{O}_{\mathbb{CP}^1}(a)) \to \mathbb{CP}^1$, which is the desired conclusion.

Next, by Lemma 2.2 of [31] we can find a suitable constant t_0 such that $2\pi t_0 c_1(\mathcal{H}_a) + f^*(2\pi c_1(\mathbb{CP}^1))$ is a Kähler class on $\mathcal{H}_a$. Then by Theorem 1.1 the continuity method (1.1) starting from any Kähler metric $\omega_0 \in 2\pi t_0 c_1(\mathcal{H}_a) + f^*(2\pi c_1(\mathbb{CP}^1))$ converges to a Kähler metric $\bar{\omega} \in 2\pi c_1(\mathbb{CP}^1)$ on $\mathbb{CP}^1$ in Gromov-Hausdorff topology. By Example 1.2 the continuity method restarting from $\bar{\omega}$ on $\mathbb{CP}^1$ will converge to a point

in Gromov-Hausdorff topology. Moreover, if we deform $\bar{\omega}$ by the normalized version (1.4)*, then it will converge in $C^\infty(X,\omega_0)$-topology to $2\omega_{FS}$, where ω_{FS} is the Fubini-Study metric on $\mathbb{CP}^1$ with $Ric(\omega_{FS}) = 2\omega_{FS}$.*

Remark 1.4. The finite time collapsing of the Kähler-Ricci flow has been studied in many papers, see [9, 10, 12, 19, 20, 21, 25, 26, 27, 31] and the references therein. For example, the collapsing of Kähler-Ricci flow on Hirzebruch surfaces is studied in [25] (also see [9] for certain generalizations), where the similar picture as in Example 1.3 is obtained by assuming certain symmetry condition on initial metrics. From the view point of Analytic Minimal Model Program with Ricci flow (in particular, see Conjecture 6.6 of [24]), the result similar to Theorem 1.1 should be true for Kähler-Ricci flow.

The rest of this paper is organized as follows. We will construct the limiting metric $\bar{\omega}_B$ on B in Section 2. Then we prove a weak convergence in Section 3 and Theorem 1.1 in Section 4. In section 5, we discuss briefly the local convergence of continuity method on Fano fibrations.

2. Construction of limiting metrics

From now on we will fix a Kähler metric $\omega_0 \in \alpha$ and a Kähler metric $\omega_B \in \beta$. By (1.2) there exists a smooth positive volume form Ω on X with

$$\sqrt{-1}\partial\bar{\partial}\log\Omega = (f^*\omega_B - \omega_0). \tag{2.1}$$

Using (1.3) and $\partial\bar{\partial}$-lemma one can choose a $\rho_b \in C^\infty(X_b,\mathbb{R})$ with

$$Ric(\omega_0|_{X_b}) - \omega_0|_{X_b} = \sqrt{-1}\partial\bar{\partial}\rho_b \tag{2.2}$$

and

$$\int_{X_b} e^{\rho_b}(\omega_0|_{X_b})^{n-k} = \int_{X_b}(\omega_0|_{X_b})^{n-k}. \tag{2.3}$$

Define a function ρ on X by setting $\rho(b,\cdot) := \rho_b(\cdot)$. Then $\rho \in C^\infty(X,\mathbb{R})$. Define

$$G = e^{-\rho}\frac{\Omega}{C_k^n\omega_0^{n-k}\wedge(f^*\omega_B)^k}.$$

Then G can be seen as a smooth positive function on B. In fact if we let ∂_f be the restriction of ∂ to the fiber direction, then by direct compution

$$\begin{aligned}\partial_f\bar{\partial}_f\log G &= -\partial_f\bar{\partial}_f\rho_b + \partial_f\bar{\partial}_f\log\Omega + Ric(\omega_0|_{X_b})\\ &= -\partial_f\bar{\partial}_f\rho_b - \omega_0|_{X_b} + Ric(\omega_0|_{X_b})\\ &= 0\end{aligned}$$

by the definition of ρ. Hence G is a constant along each fiber and descents to a smooth positive function on B.

Consider the following complex Monge-Ampère equation on B

$$(\omega_B + \sqrt{-1}\partial\bar{\partial}\psi)^k = e^\psi G\omega_B^k. \tag{2.4}$$

By Aubin-Yau theorem [1, 32] there exists a unique smooth solution ψ to (2.4) such that $\bar{\omega}_B := \omega_B + \sqrt{-1}\partial\bar{\partial}\psi$ is a Kähler metric on B. An easy observation is the following

Proposition 2.1. *The metric $\bar{\omega}_B$ defined above does not depend on the choice $\omega_B \in \beta$.*

Proof. We choose another Kähler metric $\omega'_B = \omega_B + \sqrt{-1}\partial\bar{\partial}v$ and then have the following equation similar to (2.4):

$$(\omega'_B + \sqrt{-1}\partial\bar{\partial}\psi')^k = e^{\psi'}G'\omega_B'^k, \tag{2.5}$$

where

$$G' = e^{-\rho}\frac{e^v\Omega}{C_k^n\omega_0^{n-k}\wedge(f^*\omega_B')^k}.$$

Note that

$$\begin{aligned} G'\omega_B'^k &= e^{-\rho}\frac{e^v\Omega}{C_k^n\omega_0^{n-k}\wedge(f^*\omega_B')^k}\omega_B'^k \\ &= e^{-\rho}\frac{e^v\Omega}{C_k^n\omega_0^{n-k}\wedge(f^*\omega_B')^k}\cdot\frac{\omega_0^{n-k}\wedge(f^*\omega_B')^k}{\omega_0^{n-k}\wedge(f^*\omega_B)^k}\cdot\omega_B^k \\ &= e^{-\rho}\frac{e^v\Omega}{C_k^n\omega_0^{n-k}\wedge(f^*\omega_B)^k}\cdot\omega_B^k. \end{aligned}$$

Thus, (2.5) can be translated to the following

$$(\omega_B+\sqrt{-1}\partial\bar\partial(v+\psi'))^k = e^{v+\psi'}G\omega_B^k. \tag{2.6}$$

By comparing (2.4) and (2.6), we obtain by Calabi's uniqueness theorem [5] that

$$\bar\omega_B' := \omega_B+\sqrt{-1}\partial\bar\partial(v+\psi') = \omega_B+\sqrt{-1}\partial\bar\partial\psi = \bar\omega_B.$$

Proposition 2.1 is proved. □

We will see that $\bar\omega_B$ is the unique limiting metric of the continuity method (1.1) starting from ω_0.

On the other hand, we remark that the $\bar\omega_B$ defined above does depend on the chosen metric ω_0. We shall explain how $\bar\omega_B$ changes if we choose another metric in the class α. To this end, we choose another Kähler metric $\omega_0' = \omega_0+\sqrt{-1}\partial\bar\partial u$ for some fixed $u\in C^\infty(X,\mathbb{R})$. Then we get the following equation similar to (2.4):

$$(\omega_B+\sqrt{-1}\partial\bar\partial\psi'')^k = e^{\psi''}G''\omega_B^k, \tag{2.7}$$

where

$$G'' = e^{-\rho'}\frac{e^{-u}\Omega}{C_k^n\omega_0'^{n-k}\wedge(f^*\omega_B)^k}, \tag{2.8}$$

ρ' satisfies

$$Ric(\omega_0'|_{X_b}) - \omega_0'|_{X_b} = \sqrt{-1}\partial\bar\partial\rho_b' \tag{2.9}$$

and

$$\int_{X_b} e^{\rho_b'}(\omega_0'|_{X_b})^{n-k} = \int_{X_b}(\omega_0'|_{X_b})^{n-k}. \tag{2.10}$$

Now by substituting (2.2) into (2.9), we get

$$Ric(\omega_0'|_{X_b}) = Ric(\omega_0|_{X_b}) + \sqrt{-1}\partial\bar\partial(u_b+\rho_b'-\rho_b).$$

In the above equality, $u_b(\cdot) := u(b,\cdot)$ obviously. So, there exists a constant $c(b)$ depending on b such that

$$e^{\rho_b}(\omega_0|_{X_b})^{n-k} = c_b e^{u_b+\rho_b'}(\omega_0'|_{X_b})^{n-k}. \tag{2.11}$$

We see that $c = c(b) = \frac{\int_{X_b} e^{\rho_b}(\omega_0|_{X_b})^{n-k}}{\int_{X_b} e^{u_b+\rho_b'}(\omega_0'|_{X_b})^{n-k}}$ is a smooth positive function on B.

Substituting (2.11) into (2.8) gives

$$G'' = c\cdot G$$

and hence (2.7) becomes

$$(\omega_B+\sqrt{-1}\partial\bar\partial\psi'')^k = e^{\psi''}c\cdot G\omega_B^k. \tag{2.12}$$

The solution of (2.12) will be different from that of (2.4), unless $c\equiv 1$.

Consider a very special case as follows.

Example 2.2. *Let $X = B \times F$, where F is a fixed Fano manifold, and $\omega_0 = f^*\tilde{\omega}_B + \pi^*\omega_F$, where $\tilde{\omega}_B$ (resp., ω_F) is a Kähler metric on B (resp., F) and π is the projection to F. Then $\bar{\omega}_B$ defined above satisfies*

$$Ric(\bar{\omega}_B) = -\bar{\omega}_B + \tilde{\omega}_B,$$

from which we see that $\bar{\omega}_B$ really depends on the components of ω_0 in the base directions.

3. Estimates and weak convergence

From now on, we will use the following simple reparametrization of (1.1), which will be more convenient for later discussions,

$$\begin{cases} \omega(t) = \omega_0 - (1 - e^{-t})Ric(\omega(t)) \\ \omega(0) = \omega_0. \end{cases} \tag{3.1}$$

Obviously, (3.1) has a unique solution $\omega(t)$ for $t \in [0, \infty)$ and the finite time collapsing of (1.1) at T is exactly the infinite time collapsing of (3.1).

First of all we reduce the continuity method (3.1) to a complex Monge-Ampère equation as follows. Define

$$\omega_t = e^{-t}\omega_0 + (1 - e^{-t})f^*\omega_B.$$

Then $\omega(t) = \omega_t + \sqrt{-1}\partial\bar{\partial}\varphi(t)$ solves (3.1) if $\varphi(t)$ solves

$$\begin{cases} (\omega_t + \sqrt{-1}\partial\bar{\partial}\varphi(t))^n = e^{-(n-k)t}e^{\frac{\varphi(t)}{1-e^{-t}}}\Omega \\ \varphi(0) = 0. \end{cases} \tag{3.2}$$

Lemma 3.1. *There exists a positive constant C such that for all $t \in [1, \infty)$,*

1. $\|\varphi(t)\|_{C^0(X)} \leq C$.
2. $C^{-1}e^{-(n-k)t}\Omega \leq \omega(t)^n \leq Ce^{-(n-k)t}\Omega$.
3. $tr_{\omega(t)}f^*\omega_B \leq C$.
4. $e^t|\varphi - \bar{\varphi}| \leq C$, *where* $\bar{\varphi}(b) = \frac{\int_{X_b}\varphi(\omega_0|_{X_b})^{n-k}}{\int_{X_b}(\omega_0|_{X_b})^{n-k}}$.

Proof. Note that $\omega_0^{n-k} \wedge (f^*\omega_B)^k$ is a smooth positive volume form on X, which in particular implies that there exists a positive constant C such that for all $t \in [1, \infty)$,

$$C^{-1}e^{-(n-k)t}\Omega \leq \omega_t^n \leq Ce^{-(n-k)t}\Omega. \tag{3.3}$$

Then (1), which is equivalent to (2), can be easily proved by applying the maximum principle in (3.2). The (3) can be checked by applying the Schwartz Lemma argument to $Q = \log tr_{\omega(t)}f^*\omega_B - A\varphi(t)$ (A is a large constant) since $\omega(t)$ has a uniform Ricci lower bound for $t \in [1, \infty)$. As for (4), see Lemma 2.3 of [10] for a proof. □

Lemma 3.2. *There exists a positive constant C such that for all $t \in [1, \infty)$*

$$C^{-1}\omega_t \leq \omega(t) \leq C\omega_t.$$

Proof. We give a proof by modifying method in [22] (see also [10, 11, 28]). We begin with the following inequality for $t \in [1, \infty)$, which can be checked by a direct computation as Ricci curvature of $\omega(t)$ is uniformly bounded from below for $t \in [1, \infty)$:

$$\Delta_\omega \log tr_\omega(e^{-t}\omega_0) \geq -Ctr_\omega\omega_0 - C, \tag{3.4}$$

where the positive constant C is uniform for $t \in [1, \infty)$. Moreover, similar to [22], we can find a uniform positive constant C such that for all $t \in [1, \infty)$,

$$\Delta_\omega(e^t(\varphi - \bar{\varphi})) \leq Ce^t - tr_\omega\omega_0. \tag{3.5}$$

By combining (3.4) and (3.5), we can choose a sufficiently large constant A such that for all $t \in [1, \infty)$,

$$\Delta_\omega(\log tr_\omega(e^{-t}\omega_0) - Ae^t(\varphi - \bar{\varphi})) \geq tr_\omega\omega_0 - C(A+1)e^t. \tag{3.6}$$

Now we can apply the maximum principle to (3.6) to conclude that there exists a positive constant C such that for all $t \in [1, \infty)$,

$$tr_\omega(e^{-t}\omega_0) \leq C. \tag{3.7}$$

Combining part (3) of Lemma 3.1 and (3.7), we obtain

$$tr_\omega\omega_t \leq C. \tag{3.8}$$

Moreover,

$$\begin{aligned} tr_{\omega_t}\omega &\leq \frac{1}{(n-1)!}(tr_\omega\omega_t)^{n-1}\frac{\omega^n}{\omega_t^n} \\ &\leq C. \end{aligned} \tag{3.9}$$

We conclude (3.4) from (3.8) and (3.9). □

A direct consequence of Lemma 3.2 is the following

Corollary 3.3. *There exists a positive constant C such that for all $b \in B$ and $t \in [1, \infty)$*

$$C^{-1}e^{-\frac{t}{2}} \leq diam(X_b, \omega(t)|_{X_b}) \leq Ce^{-\frac{t}{2}}.$$

Combining Lemma 3.1 (1) and Lemma 3.2 we obtain

Proposition 3.4. *For any $\theta \in (0,1)$, there exists a positive constant C_θ such that for all $t \in [1, \infty)$*

$$\|\varphi(t)\|_{C^{1,\theta}(X,\omega_0)} \leq C_\theta.$$

Now we show a weak convergence of $\omega(t)$ to $f^*\bar{\omega}_B$.

Proposition 3.5. *As $t \to \infty$, $\omega(t) \to f^*\bar{\omega}_B$ as currents on X and the convergence takes place in $C^{1,\theta}$-topology at the level of Kähler potentials for any $\theta \in (0,1)$.*

Proof. By Proposition 3.4, for any fixed $\theta \in (0,1)$ we choose a time sequence t_j increasing to ∞ such that $\varphi(t_j) \to \varphi_\infty$ in $C^{1,\theta}(X, \omega_0)$-topology for some $\varphi_\infty \in C^{1,\alpha}(X, \omega_0)$. In particular $\omega(t_j) \to f^*\omega_B + \sqrt{-1}\partial\bar{\partial}\varphi_\infty$ as currents on X and $f^*\omega_B + \sqrt{-1}\partial\bar{\partial}\varphi_\infty$ is positive as current. For any $b \in B$, by restricting $f^*\omega_B + \sqrt{-1}\partial\bar{\partial}\varphi_\infty$ to X_b we see that $\sqrt{-1}\partial\bar{\partial}\varphi_\infty|_{X_b} \geq 0$ and hence φ_∞ is constant on every fiber X_b. Therefore φ_∞ descents to a function, still denote it by φ_∞, on B with $C^{1,\theta}(B, \omega_B)$-regularity. We want to show that φ_∞ is a weak solution of (2.4), namely, for any $\eta \in C^\infty(B, \mathbb{R})$ φ_∞ satisfies

$$\int_B \eta(\omega_B + \sqrt{-1}\partial\bar{\partial}\varphi_\infty)^k = \int_B \eta e^{\varphi_\infty} G\omega_B^k. \tag{3.10}$$

The following arguments are similar to [28]. To begin with, using the equation (3.2) we have

$$\int_X f^*\eta e^{(n-k)t}(\omega_t + \sqrt{-1}\partial\bar{\partial}\varphi(t))^n = \int_X f^*\eta e^{\frac{\varphi(t)}{1-e^{-t}}}\Omega. \tag{3.11}$$

On the one hand, for the right hand side of (3.11),

$$\begin{aligned}\int_X f^*\eta e^{\frac{\varphi(t_j)}{1-e^{-t_j}}}\Omega &\to \int_X (f^*\eta)e^{\varphi_\infty}\Omega\\ &= \int_X (f^*\eta)e^{\varphi_\infty}(C_k^n G e^{\rho}\omega_0^{n-k}\wedge(f^*\omega_B)^k)\\ &= \left(\int_{b\in B}\eta e^{\varphi_\infty}G\omega_B^k\right)\left(\int_{X_b}C_k^n e^{\rho_b}(\omega_0|_{X_b})^{n-k}\right) \qquad (3.12)\end{aligned}$$

On the other hand, for the left hand side of (3.11),

$$\begin{aligned}&\int_X f^*\eta e^{(n-k)t}(\omega_t+\sqrt{-1}\partial\bar\partial\varphi(t))^n\\ &= \int_X f^*\eta e^{(n-k)t}\left(((1-e^{-t})f^*\omega_B+\sqrt{-1}\partial\bar\partial\bar\varphi)+(e^{-t}\omega_0+\sqrt{-1}\partial\bar\partial(\varphi-\bar\varphi))\right)^n\\ &= \int_X f^*\eta e^{(n-k)t}\sum_{i=0}^{k}C_i^n((1-e^{-t})f^*\omega_B+\sqrt{-1}\partial\bar\partial\bar\varphi)^i\wedge(e^{-t}\omega_0+\sqrt{-1}\partial\bar\partial(\varphi-\bar\varphi))^{n-i}\\ &= \int_X f^*\eta e^{(n-k)t}\sum_{i=0}^{k}\sum_{l=0}^{n-i}C_i^n C_l^{n-i}((1-e^{-t})f^*\omega_B+\sqrt{-1}\partial\bar\partial\bar\varphi)^i\wedge(e^{-t}\omega_0)^{n-i-l}\wedge(\sqrt{-1}\partial\bar\partial(\varphi-\bar\varphi))^l\\ &=: \sum_{i=0}^{k}\sum_{l=0}^{n-i}A_{i,l},\end{aligned}$$

where

$$A_{i,l}=\int_X f^*\eta e^{(n-k)t}C_i^nC_l^{n-i}((1-e^{-t})f^*\omega_B+\sqrt{-1}\partial\bar\partial\bar\varphi)^i\wedge(e^{-t}\omega_0)^{n-i-l}\wedge(\sqrt{-1}\partial\bar\partial(\varphi-\bar\varphi))^l$$

and $0\le i\le k$ and $0\le l\le n-i$.

(1) If $i<k$ and $l=0$, then $A_{i,l}\le Ce^{-(k-i)t}$.

(2) If $i<k$ and $l>0$, then by integration by part we have

$$A_{i,l}=\int_X e^{(n-k)t}C_i^nC_l^{n-i}(\varphi-\bar\varphi)\sqrt{-1}\partial\bar\partial f^*\eta\wedge((1-e^{-t})f^*\omega_B+\sqrt{-1}\partial\bar\partial\bar\varphi)^i\wedge(e^{-t}\omega_0)^{n-i-l}\wedge(\sqrt{-1}\partial\bar\partial(\varphi-\bar\varphi))^{l-1} \qquad (3.13)$$

By (3.4) we know

$$-Ce^{-t}\le\sqrt{-1}\partial\bar\partial(\varphi-\bar\varphi)|_{X_b}=\sqrt{-1}\partial\bar\partial\varphi|_{X_b}\le Ce^{-t} \qquad (3.14)$$

Combining with part (4) of Lemma 3.1, we conclude that

$$A_{i,l}\le Ce^{(n-k)t}e^{-t}e^{-(n-i-l)t}e^{-((l-1)-(k-i-1))t}=Ce^{-t} \qquad (3.15)$$

(3) If $i=k$ and $l>0$, then by integration by part we see that $A_{i,l}=0$.

(4) If $i=k$ and $l=0$, then

$$\begin{aligned}A_{k,0}&=\int_X f^*\eta e^{(n-k)t}C_k^n((1-e^{-t})f^*\omega_B+\sqrt{-1}\partial\bar\partial\bar\varphi)^k\wedge(e^{-t}\omega_0)^{n-k}\\ &=\int_X f^*\eta C_k^n((1-e^{-t})f^*\omega_B+\sqrt{-1}\partial\bar\partial\bar\varphi)^k\wedge\omega_0^{n-k}\\ &\to\left(\int_{b\in B}\eta(\omega_B+\sqrt{-1}\partial\bar\partial\varphi_\infty)^k\right)\left(C_k^n\int_{X_b}(\omega_0|_{X_b})^{n-k}\right),\end{aligned}$$

where the convergence in last line is taken along $t_j\to\infty$. Combining the above cases (1)-(4), we conclude that the left hand side of (3.11) converges to

$$\left(\int_{b\in B}\eta(\omega_B+\sqrt{-1}\partial\bar\partial\varphi_\infty)^k\right)\left(C_k^n\int_{X_b}(\omega_0|_{X_b})^{n-k}\right) \qquad (3.16)$$

as $t_j \to \infty$. Therefore, by the normalization (2.3) and the fact that $\int_{X_b}(\omega_0|_{X_b})^{n-k}$ is a positive constant independent in b, we get the desired equation (3.10). In particular, φ_∞ is a continuous solution to (2.4). By applying the comparison principle of complex Monge-Ampère equations (see Theorem 2.1 in [14]), we have

$$\begin{aligned}\int_{\{\varphi_\infty \le \psi\}} e^{\psi} G\omega_B^k &= \int_{\{\varphi_\infty \le \psi\}} (\omega_B + \sqrt{-1}\partial\bar\partial\psi)^k \\ &\le \int_{\{\varphi_\infty \le \psi\}} (\omega_B + \sqrt{-1}\partial\bar\partial\varphi_\infty)^k \\ &= \int_{\{\varphi_\infty \le \psi\}} e^{\varphi_\infty} G\omega_B^k,\end{aligned}$$

which implies

$$\varphi_\infty \ge \psi \tag{3.17}$$

on B. Similarly, we have

$$\varphi_\infty \le \psi \tag{3.18}$$

on B. Combining (3.17) and (3.18) gives

$$\varphi_\infty = \psi$$

and hence $\varphi(t) \to \psi$ in $C^{1,\theta}$-topology without passing to a subsequence. Consequently, as $t \to \infty$, $\omega(t) \to \pi^*\omega_B$ in the current sense. The proof of Proposition 3.5 is completed. □

4. Proof of Theorem 1.1

In this section, we will give a proof of Theorem 1.1 by using the strategy developed in [29]. To this end, we need more estimates. Easily, (3.2) can be rewritten as follows

$$\varphi(t) = (1 - e^{-t}) \log \frac{e^{(n-k)t}(\omega_t + \sqrt{-1}\partial\bar\partial\varphi(t))^n}{\Omega}$$

A direct computation gives

$$\partial_t\varphi = e^{-t} \log \frac{e^{(n-k)t}(\omega_t + \sqrt{-1}\partial\bar\partial\varphi(t))^n}{\Omega} + (1-e^{-t})(n-k+tr_\omega(f^*\omega_B - \omega_t) + \Delta_\omega(\partial_t\varphi)) \tag{4.1}$$

and

$$\begin{aligned}\partial_t\partial_t\varphi =& (1 - e^{-t})\Delta_\omega(\partial_t\partial_t\varphi) - e^{-t} \log \frac{e^{(n-k)t}(\omega_t + \sqrt{-1}\partial\bar\partial\varphi(t))^n}{\Omega} \\ &+ (1 - 3e^{-t})tr_\omega(\omega_t - f^*\omega_B) + 2e^{-t}(n-k) - (1-e^{-t})|\partial_t\omega|^2_\omega. \end{aligned}\tag{4.2}$$

By applying the maximum principle in (4.1) we see that there exists a positive constant C such that for all $t \in 1$,

$$\|\partial_t\varphi\|_{C^0(X)} \le C \tag{4.3}$$

and hence

$$\|\Delta_\omega\partial_t\varphi\|_{C^0(X)} \le C. \tag{4.4}$$

Using (4.4) and the maximum principle in (4.2) gives

$$\sup_{X\times[1,\infty)} \partial_t\partial_t\varphi \le C \tag{4.5}$$

for some positive constant C.

With (4.3) and (4.5), we can apply the arguments in [29] to prove the following analogue of Lemma 4.6 of [29].

Lemma 4.1. *There exists a positive function $H(t)$ with $H(t) \to 0$ as $t \to \infty$ such that*

$$\|\varphi - f^*\psi\|_{C^0(X)} \leq H(t) \tag{4.6}$$

and

$$\|\partial_t \varphi\|_{C^0(X)} \leq H(t). \tag{4.7}$$

Consequently,

$$\|\varphi + \partial_t \varphi - f^*\psi\|_{C^0(X)} \leq 2H(t). \tag{4.8}$$

Proof. For the sake of convenience, we present a proof here. Firstly, we set $H_1(t) := \|\varphi(t) - f^*\psi\|_{C^0(X)}$, which by Proposition 3.5 goes to zero as $t \to \infty$. For (4.7), we argue by contradiction. Assume there exists a constant $\delta_0 > 0$, $x_i \in X$ and $t_i \to \infty$ such that

$$\partial_t \varphi(t_i, x_i) \leq -\delta_0.$$

Then by using (4.5) we have

$$\partial_t \varphi(t, x_i) \leq -\frac{\delta_0}{2} \tag{4.9}$$

for all $t \in [t_i, t_i + \frac{\delta_0}{2C}]$. Integrating (4.9) over t gives

$$(\varphi - f^*\psi)(t_i + \frac{\delta_0}{2C}, x_i) \leq (\varphi - f^*\psi)(t_i, x_i) - \frac{\delta_0^2}{4C},$$

which implies

$$-H_1(t_i + \frac{\delta_0}{2C}) \leq H_1(t_i) - \frac{\delta_0^2}{4C}.$$

Let $i \to \infty$ we get a contradiction. Therefore, we conclude that

$$\liminf_{t \geq 1} \inf_{t \geq 1} \partial_t \varphi \geq 0. \tag{4.10}$$

Similarly, we have

$$\limsup_{t \geq 1} \sup_{t \geq 1} \partial_t \varphi \leq 0. \tag{4.11}$$

Now (4.7) follows from (4.10) and (4.11). Finally, (4.6) and (4.7) imply (4.8) immediately. □

Then we have the following

Lemma 4.2. *There exist a positive constant C such that for all $t \geq 1$,*

$$\sup_X (tr_{\omega(t)} f^*\bar{\omega}_B - k) \leq C\sqrt{H(t)}. \tag{4.12}$$

Proof. This lemma follows from a maximum principle argument as in [29]. For completeness we present a proof here. To begin with, using (4.1), we have

$$\begin{aligned}\Delta_\omega(\varphi + \partial_t\varphi - f^*\psi) &= -(tr_\omega f^*\bar{\omega}_B - k) + \frac{\partial_t\varphi}{1 - e^{-t}} - \frac{e^{-t}}{1 - e^{-t}} \log \frac{e^{(n-k)t}(\omega_t + \sqrt{-1}\partial\bar{\partial}\varphi(t))^n}{\Omega} \\ &\leq -(tr_\omega f^*\bar{\omega}_B - k) + CH(t) + Ce^{-t}, \end{aligned} \tag{4.13}$$

where we have used (4.7) and part (2) of Lemma 3.1 in the last inequality. On the other hand, by a Schwarz Lemma argument (see, e.g., [22]) we can choose a positive constant C such that for all $t \geq 1$,

$$\Delta_\omega tr_\omega f^*\bar{\omega}_B \geq -\frac{1}{1 - e^{-1}} tr_\omega f^*\bar{\omega}_B - C(tr_\omega f^*\bar{\omega}_B)^2, \tag{4.14}$$

Moreover, using part (3) of Lemma 3.1 and the fact that ω_B is equivalent to $\bar{\omega}_B$ we have

$$\Delta_\omega tr_\omega f^*\bar{\omega}_B \geq -C. \tag{4.15}$$

Combining (4.13) and (4.15) we get

$$\Delta_\omega\left((tr_\omega f^*\bar{\omega}_B - k) - \frac{(\varphi + \partial_t\varphi - f^*\psi)}{\sqrt{H(t)}}\right) \geq \frac{tr_\omega f^*\bar{\omega}_B - k}{\sqrt{H(t)}} - C\sqrt{H(t)} - \frac{e^{-t}}{\sqrt{H(t)}} - C.$$

Note that we may assume $\sqrt{H(t)} \geq Ce^{-t}$ for all $t \in [1, \infty)$. Thus we arrive at

$$\Delta_\omega\left((tr_\omega f^*\bar{\omega}_B - k) - \frac{(\varphi + \partial_t\varphi - f^*\psi)}{\sqrt{H(t)}}\right) \geq \frac{tr_\omega f^*\bar{\omega}_B - k}{\sqrt{H(t)}} - C. \tag{4.16}$$

Now by applying the maximum principle in (4.16) and using (4.8), the proof of (4.12) is completed. □

Now we determine the limiting metrics of the normalized metric along the fibers. Motivated by an observation in Example 1.2, these limiting metrics should be given by the follow equations (4.17). For any $b \in B$, consider

$$\begin{cases} (\omega_0|_{X_b} + \sqrt{-1}\partial\bar{\partial}W_b)^{n-k} = e^{\rho_b}(\omega_0|_{X_b})^{n-k} \\ \displaystyle\int_{X_b} W_b(\omega_0|_{X_b})^{n-k} = 0. \end{cases} \tag{4.17}$$

By Yau's theorem [32] we know (4.17) admits a unique smooth solution w_b such that $\omega_0|_{X_b} + \sqrt{-1}\partial\bar{\partial}W_b$ is a Kähler metric on X_b. If we define $W(b, \cdot) := W_b(\cdot)$, then $W \in C^\infty(X, \mathbb{R})$. Set $\bar{\omega}_0 := \omega_0 + \sqrt{-1}\partial\bar{\partial}W$ and $\bar{\omega}_{0,b} := \omega_0|_{X_b} + \sqrt{-1}\partial\bar{\partial}W_b = \bar{\omega}_0|_{X_b}$.

Remark 4.3. Note that by definition $Ric(\bar{\omega}_{0,b}) = \omega_0|_{X_b}$.

The following proposition can be regarded as a family version of some discussions in Example 1.2.

Proposition 4.4. *For every $b \in B$ we have, as $t \to \infty$,*

$$e^t\omega|_{X_b} \to \bar{\omega}_{0,b} \tag{4.18}$$

in $C^\infty(X_b, \omega_0|_{X_b})$-topology.

Proof. Using the equation (3.2) we have

$$\begin{aligned}
(\omega_0|_{X_b} + \sqrt{-1}\partial\bar{\partial}(e^t\varphi|_{X_b}))^{n-k} &= (e^t\omega|_{X_b})^{n-k} \\
&= \frac{(e^t\omega|_{X_b})^{n-k}}{(\omega_0|_{X_b})^{n-k}}(\omega_0|_{X_b})^{n-k} \\
&= \frac{(e^t\omega)^{n-k} \wedge (f^*\bar{\omega}_B)^k}{(\omega_0)^{n-k} \wedge (f^*\bar{\omega}_B)^k}(\omega_0|_{X_b})^{n-k} \\
&= \frac{\omega^{n-k} \wedge (f^*\bar{\omega}_B)^k}{\omega^n} \frac{e^{(n-k)t}\omega^n}{\omega_0^{n-k} \wedge (f^*\bar{\omega}_B)^k}(\omega_0|_{X_b})^{n-k} \\
&= \frac{C_k^n\omega^{n-k} \wedge (f^*\bar{\omega}_B)^k}{\omega^n} \frac{e^{\frac{\varphi}{1-e^{-t}}}\Omega}{C_k^n\omega_0^{n-k} \wedge (f^*\bar{\omega}_B)^k}(\omega_0|_{X_b})^{n-k} \\
&= e^{\frac{\varphi}{1-e^{-t}} - f^*\psi} \cdot \frac{C_k^n\omega^{n-k} \wedge (f^*\bar{\omega}_B)^k}{\omega^n} e^\rho(\omega_0|_{X_b})^{n-k}.
\end{aligned} \tag{4.19}$$

Set

$$h = e^{\frac{\varphi}{1-e^{-t}} - f^*\psi} \cdot \frac{C_k^n\omega^{n-k} \wedge (f^*\bar{\omega}_B)^k}{\omega^n}$$

and

$$h_b = h|_{X_b}.$$

By Proposition 3.5 we know

$$e^{\frac{\varphi}{1-e^{-t}}-f^*\psi}\to 1$$

in $C^0(X)$-topology. On the other hand, we have

$$\frac{C_k^n\omega^{n-k}\wedge(f^*\bar{\omega}_B)^k}{\omega^n}\leq\left(\frac{tr_\omega f^*\bar{\omega}_B}{k}\right)^k.$$

Hence, firstly by Yau's C^0-estimate we know $e^t(\varphi-\bar{\varphi})|_{X_b}$ is uniformly bounded. Moreover, combining with Lemma 4.2, we can find a positive function $K_0(t)$ with $K_0(t)\to 0$ as $t\to\infty$ such that for all $b\in B$ and $t\in[1,\infty)$,

$$h_b-1\leq K_0(t) \tag{4.20}$$

and hence

$$(h_b-1)^+\leq K_0(t), \tag{4.21}$$

where $(h_b-1)^+:=\max\{h_b-1,0\}$ and $(h_b-1)^-:=\max\{-(h_b-1),0\}$. Note that

$$0=\int_{X_b}(h_b-1)e^\rho(\omega_0|_{X_b})^{n-k}=\int_{X_b}(h_b-1)^+e^\rho(\omega_0|_{X_b})^{n-k}-\int_{X_b}(h_b-1)^-e^\rho(\omega_0|_{X_b})^{n-k}.$$

So, by using (4.21) we have

$$\int_{X_b}(h_b-1)^-e^\rho(\omega_0|_{X_b})^{n-k}=\int_{X_b}(h_b-1)^+e^\rho(\omega_0|_{X_b})^{n-k}\leq K_0(t)\int_{X_b}e^\rho(\omega_0|_{X_b})^{n-k}=CK_0(t)$$

and hence

$$\int_{X_b}|h_b-1|e^\rho(\omega_0|_{X_b})^{n-k}=\int_{X_b}(h_b-1)^+e^\rho(\omega_0|_{X_b})^{n-k}+\int_{X_b}(h_b-1)^-e^\rho(\omega_0|_{X_b})^{n-k}\leq CK_0(t).$$

Now by applying Kołodziej's stability theorem [14] we can choose a unique bounded function $C_b(t)$, which depends on $b\in B$ and $t\in[1,\infty)$ continuously and is constant along fiber X_b, such that

$$\|e^t(\varphi-\bar{\varphi})|_{X_b}-C_b-W_b\|_{L^\infty(X_b)}\leq CK_0^{\frac{1}{n-k+4}}(t),$$

Immediately, (4.18) holds as currents on X_b. Moreover, for any $k\in\mathbb{N}$, we can obtain the k^{th} order estimate of $e^t\omega|_{X_b}$ with respect to $\omega_0|_{X_b}$ by similar arguments in [30], which is uniform in $b\in B$. The proof of Proposition 4.4 is completed. □

We also need the following

Lemma 4.5. *There exists a positive function $H_1(t)$ with $H_1(t)\to 0$ as $t\to\infty$ such that for all $t\in[1,\infty)$,*

$$\|e^t\omega|_{X_b}-\bar{\omega}_{0,b}\|_{C^0(X_b,\omega_0|_{X_b})}\leq H_1(t). \tag{4.22}$$

Proof. Firstly, the uniform first order estimate of $e^t\omega|_{X_b}$ with respect to $\omega_0|_{X_b}$ implies that $\|\nabla h_b\|_{\omega_0|_{X_b}}$ is uniformly bounded for $b\in B$. Hence we have

$$\|h_b-1\|_{C^0(X_b)}\leq K_1(t)$$

for some K_1 that goes to 0 as $t\to\infty$, which is equivalent to

$$\left\|\frac{(e^t\omega|_{X_b})^{n-k}}{(\bar{\omega}_{0,b})^{n-k}}-1\right\|_{C^0(X_b)}\leq K_1(t). \tag{4.23}$$

Moreover, if we set

$$\tilde{h}(b)=\frac{(e^t\omega|_{X_b})\wedge\bar{\omega}_{0,b}^{n-k-1}}{\bar{\omega}_{0,b}^{n-k}},$$

then by discussions in Proposition 4.4 we know $\|\nabla\tilde{h}(b)\|_{\omega_0|_{X_b}}$ is uniformly bounded for $b \in B$ and for any $b \in B$,

$$\int_{X_b} (\tilde{h}(b) - 1)\bar{\omega}_{0,b} = 0.$$

Also, note that

$$\begin{aligned}\tilde{h}(b) &= \frac{1}{n-k} tr_{\bar{\omega}_{0,b}}(e^t\omega_0|_{X_b}) \\ &\geq \left(\frac{(e^t\omega|_{X_b})^{n-k}}{(\bar{\omega}_{0,b})^{n-k}}\right)^{\frac{1}{n-k}} \\ &\geq 1 - K_2(t)\end{aligned}$$

by (4.23). Here $K_2(t)$ is a positive function that goes to 0 as $t \to \infty$. Now we can conclude from Lemma 2.4 of [29] that

$$\|\tilde{h}(b) - 1\|_{C^0(X_b)} \leq K_3(t) \tag{4.24}$$

for some positive function $K_3(t)$ that goes to 0 as $t \to \infty$. With (4.23) and (4.24), we conclude (4.22) immediately by applying Lemma 2.6 of [29]. □

Remark 4.6. From the arguments in Lemma 4.5 we see that Proposition 4.4 can be proved in a more elementary way without involving Kołodziej's stability theorem.

With Lemma 4.2, Proposition 4.4 and Lemma 4.5, we can prove the following proposition by similar arguments in [29].

Proposition 4.7. *As $t \to \infty$, $\|\omega(t) - \bar{\omega}_t\|_{C^0(X,\omega_0)} \to 0$, where $\bar{\omega}_t := e^{-t}\bar{\omega}_0 + (1 - e^{-t})f^*\bar{\omega}_B$.*

Note that $\bar{\omega}_t$ defined above is a Kähler metric on X for all sufficiently large t.

Finally, we are able to prove Theorem 1.1.

Proof of Theorem 1.1. Note that Proposition 4.7 implies $\|\omega(t) - f^*\bar{\omega}_B\|_{C^0(X,\omega_0)} \to 0$ and hence Theorem 1.1.

5. The continuity method on Fano fibrations

We end this paper by discussing how to extend the results in previous sections to a Fano fibration. Let $\hat{X}$ be an n-dimensional compact connected Kähler manifold admitting a Fano fibration over a k-dimensional compact connected Kähler manifold $\hat{B}$ with $k < n$. Namely, there exists a surjective holomorphic map $g : \hat{X} \to \hat{B}$ with connected fibers such that $K_{\hat{X}}^{-1}$ is g-ample. If we denote the critical locus of $\hat{f}$ by S, then $K_{\hat{X}}^{-1}|_{\hat{X}_b} \cong K_{\hat{X}_b}^{-1}$ is ample for all $b \in \hat{B} \setminus S$. Denote $\hat{B}_{reg} := \hat{B} \setminus S$ and $\hat{X}_{reg} := \hat{X} \setminus g^{-1}(S)$. Obviously, $g : \hat{X}_{reg} \to \hat{B}_{reg}$ is a surjective holomorphic submersion. According to Lemma 2.2 in [31], we fix a Kähler class $\hat{\alpha}$ on $\hat{X}$ and a Kähler class $\hat{\beta}$ on $\hat{B}$ such that $\hat{\alpha} = 2\pi c_1(\hat{X}) + g^*\hat{\beta}$. Also we fix a Kähler metric $\hat{\omega}_0 \in \hat{\alpha}$ and a Kähler metric $\omega_{\hat{B}} \in \hat{\beta}$.

In the obvious way, we can define $\hat{\rho} \in C^\infty(\hat{X}_{reg})$ and $\hat{G} \in C^\infty(\hat{X}_{reg})$ as in Section 2. A useful observation is the following

Lemma 5.1. *There exists a constant $p > 1$ such that $\hat{G} \in L^p(\hat{B}, \omega_{\hat{B}}^k)$.*

This lemma can be checked by the argument in Proposition 3.2 of [23] and the normalization $\int_{\hat{X}_b} e^{\hat{\rho}_b}(\hat{\omega}_0|_{\hat{X}_b})^{n-k} = \int_{\hat{X}_b}(\hat{\omega}_0|_{\hat{X}_b})^{n-k} = C$, where C is a positive constant independent in $b \in \hat{B}_{reg}$. We should remark that, when we are trying to estimate $\hat{G}$ near the singular set S, this normalization releases us from estimating $\hat{\rho}$ near singular fibers.

Next, applying the results in [13, 15, 23] gives a unique solution

$$\hat{\psi} \in C^{\hat{\theta}}(\hat{B}, \omega_{\hat{B}}) \bigcap C^{\infty}(\hat{B}_{reg}, \omega_{\hat{B}})$$

for some suitable Hölder exponent $\hat{\theta} \in (0,1)$ to the following equation:

$$(\omega_{\hat{B}} + \sqrt{-1}\partial\bar{\partial}\hat{\psi})^k = e^{\hat{\psi}}\hat{G}\omega_{\hat{B}}^k.$$

Denote $\bar{\omega}_{\hat{B}} := \omega_{\hat{B}} + \sqrt{-1}\partial\bar{\partial}\hat{\psi}$, which is a closed positive $(1,1)$-current on $\hat{B}$ and a smooth Kähler metric on $\hat{B}_{reg}$. Let $\hat{\omega}(t)$ be the unique solution of (1.1) (or (3.1)) starting from $\hat{\omega}_0$ on $\hat{X}$. Then by using the techniques developed in [22, 28, 29] and our arguments in previous sections, we have

Proposition 5.2. *For any $K \subset\subset \hat{X}_{reg}$, $\hat{\omega}(t) \to f^*\bar{\omega}_{\hat{B}}$ in $C^0(K, \hat{\omega}_0)$-topology as $t \to \infty$.*

We do expect that the metric completion of $(\hat{B}_{reg}, \bar{\omega}_{\hat{B}})$ is compact and is the unique Gromov-Hausdorff limit of $(\hat{X}, \hat{\omega}(t))$ as $t \to \infty$.

Acknowledgements

Y. Zhang is grateful to Prof. Huai-Dong Cao for constant encouragement and support. He also appreciates the encouragement from Prof. Chengjie Yu, as well as his invitation to visit Shantou University. Part of this work was carried out while Y. Zhang was visiting Capital Normal University and Shantou University, which he would like to thank for the warm hospitality.

References

[1] Aubin, T., Équations du type Monge-Ampère sur les variétés kähleriennes compactes, C. R. Acad. Sci. Paris Sér. A-B 283 (1976), no. 3, Aiii, A119-A121
[2] Aubin, T., Réduction de cas positif de léquation de Monge-Ampère sur les variétés kähleriennes compactes à la démonstration dune inégalité. Journal of Functional Analysis (2) 57 (1984), 143-153
[3] Bando, S. and Mabuchi, T., Uniqueness of Kähler-Einstein metrics modulo connected group actions, in Algebraic Geometry, Sendai, 1985 (T. Oda, Ed.), Advanced Studies in Pure Mathematics 10, Kinokuniya, 1987, 11-40.
[4] Barth, W., Hulek, K., Peters, C. and Van de Ven, A., Compact complex surfaces, Springer (2003)
[5] Calabi, E., On Kähler manifolds with vanishing canonical class, Algebraic Geometry and Topology, A Symposium in honor of S. Lefschatz, Princeton, 1955, 78-89
[6] Choi, Y.-J., Semi-positivity of fiberwise Ricci-flat metrics on Calabi-Yau fibrations, arXiv:1508.00323v3
[7] Cox, D., Little, J. and Schenck, H., Toric varieties, Graduate Studies in Mathematics, AMS, 124 (2010)
[8] Datar, V. and Székelyhidi, G., Kähler-Einstein metrics along smooth continuity method, Geom. Funct. Anal. 26 (2016) 975-1010
[9] Fong, F., Kähler-Ricci flow on projective bundles over Kähler-Einstein manifolds, Trans. Amer. Math. Soc. (2014) 366, 563-598
[10] Fong, F., On the collapsing rate of the Kähler-Ricci flow with finite time singularity, J. Geom. Anal. 25 (2015), no. 2, 1098-1107
[11] Fong, F. and Zhang, Z., The collapsing rate of The Kähler-Ricci flow with regular infinite time singularities, J. Reine Angew. Math. 703 (2015), 95-113
[12] Fu, X. and Zhang, S., The Kähler-Ricci flow on Fano bundles, arXiv:1610.03366v1
[13] Kołodziej, S., The complex Monge-Ampère equation, Acta Math. 180 (1), 69-117 (1998)
[14] Kołodziej, S., The complex Monge-Ampère equation on compact Kähler manifolds, Indiana Univ. Math. J. 52 (2003), no. 3, 667-686
[15] Kołodziej, S., Hölder continuity of solutions to the complex Monge-Ampère equation with right hand side in L^p. The case of compact Kähler manifolds, Math. Ann. 342, 379-386
[16] La Nave, G. and Tian, G., A continuity method to construct canonical metrics, Math. Ann. 365 (2016), 911-921

[17] Matsuki, K., Introduction to the Mori program, Universitext, Springer-Verlag, New York, 2002
[18] Rubinstein, Y., Some discretizations of geometric evolution equations and the Ricci iteration on the space of Kahler metrics, Adv. Math. 218 (2008), 1526-1565
[19] Šešum, N. and Tian, G., Bounding scalar curvature and diameter along the Kähler-Ricci flow (after Perelman), J. Inst. Math. Jussieu 7 (2008), 575-587
[20] Song, J., Finite time extinction of the Kähler-Ricci flow, Math. Res. Lett. 21 (2014), no. 6, 1435-1449
[21] Song, J., Székelyhidi, G. and Weinkove, B., The Kähler-Ricci flow on projective bundles, Int. Math. Res. Not. 2013, no. 2, 243-257
[22] Song, J. and Tian, G., The Kähler-Ricci flow on surfaces of positive Kodaira dimension, Invent. Math., 170, 609-653 (2006)
[23] Song, J. and Tian, G., Canonical measures and Kähler-Ricci flow, J. Amer. Math. Soc. 25 (2012), no. 2, 303-353
[24] Song, J. and Tian, G., The Kähler-Ricci flow through singularities, to appear in Invent. Math. (2016), doi:10.1007/s00222-016-0674-4
[25] Song, J. and Weinkove, B., The Kähler-Ricci flow on Hirzebruch surfaces, J. Reine Angew. Math., 659 (2011), 141-168
[26] Song, J. and Yuan, Y., Metric flips with Calabi ansatz, Geom. Funct. Anal. Vol. 22 (2012), 240-265
[27] Tian, G. and Zhu, X., Convergence of Kähler-Ricci flow, J. Amer. Math. Soc. 20, 675-699 (2007)
[28] Tosatti, V., Adiabatic limits of Ricci-flat Kähler metrics, J. Diff. Geom. 84 (2010) 427-453
[29] Tosatti, V., Weinkove, B. and Yang, X., The Kähler-Ricci flow, Ricci-flat metrics and collapsing limits, arXiv:1408.0161v1
[30] Tosatti, V. and Zhang, Y.-G., Infinite time singularities of the Kähler-Ricci flow, Geometry and Topology, 19 (2015), 2925-2948
[31] Tosatti, V. and Zhang, Y.-G., Finite time collapsing of the Kähler-Ricci flow on threefolds, to appear in Ann. Sc. Norm. Super. pisa CI. Sci., arXiv:1507.08397v2
[32] Yau, S.-T., On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, I, Comm. Pure Appl. Math. 31 (1978) 339-411

Department of mathematics, University of Macau, Macau, China
E-mail address: yashanzh@163.com

Department of mathematics, Capital Normal University, Beijing, China
E-mail address: zhleigo@aliyun.com